

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

BQ

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/18	A1	(11) International Publication Number: WO 96/20004
		(43) International Publication Date: 4 July 1996 (04.07.96)

(21) International Application Number: PCT/JP95/02707

(22) International Filing Date: 27 December 1995 (27.12.95)

(30) Priority Data:
6/337884 27 December 1994 (27.12.94) JP(71) Applicant (for all designated States except US): SNOW
BRAND MILK PRODUCTS CO., LTD. [JP/JP]; 1-1,
Naebocho 6-chome, Higashi-ku, Sapporo-shi, Hokkaido
065 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SHIOTA, Akira [JP/JP];
544-5, Shizuwa, Iwafunemachi, Shimotsuga-gun, Tochigi
329-43 (JP). FUJISE, Nobuaki [JP/JP]; Ishibashi Heights
201, 223-2, Ishibashi, Ishibashimachi, Shimotsuga-gun,
Tochigi 329-05 (JP). MASUNAGA, Hiroaki [JP/JP]; 25-9,
Ekihigashicho, Mibumachi, Shimotsuga-gun, Tochigi 321-
02 (JP). HIGASHIO, Kanji [JP/JP]; 1769-10, Yamada,
Kawagoe-shi, Saitama 350 (JP).(74) Agents: FUJINO, Seiya et al.; Mitsuhama Building, 8F, 2-1,
Yotsuya 1-chome, Shinjuku-ku, Tokyo 160 (JP).(81) Designated States: AU, CA, FI, HU, KR, NO, NZ, US,
European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR,
IE, IT, LU, MC, NL, PT, SE).

Published

*With international search report.**Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.*

(54) Title: THERAPEUTIC AGENT FOR DISORDER OF LIPID METABOLISM

(57) Abstract

The present invention relates to a therapeutic agent for treatment of disorder of lipid metabolism comprising TCF-II as an effective ingredient. The therapeutic agent of the present invention is useful for treatment of disorder of lipid metabolism caused by diabetes, endocrinological diseases, such as thyroid dysfunction and adrenal dysfunction, chronic metabolism disorder, chronic renal failure including nephrosis, uremia, cancer, dyscrasia, adverted effects of antitumor agents, gastrointestinal (including hepatic and pancreatic) impairment and so on. Hypercholesterolemia, hypolipidemia or hypo-high-density-lipoproteinemia is exemplified as a disorder of lipid metabolism. The therapeutic agent of the present invention normalizes not only hyperlipidemia but also hypolipidemia. In addition, it improves fatty liver.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Title of the Invention

THERAPEUTIC AGENT FOR DISORDER OF LIPID METABOLISM

Field of the Invention

The present invention relates to a therapeutic agent for treatment of disorder of lipid metabolism comprising TCF-II as an effective ingredient. The therapeutic agent of the present invention is useful for treatment of disorder of lipid metabolism such as hypercholesterolemia, hypolipidemia, hypo-high-density-lipoproteinemia etc., which is induced by diabetes, endocrinological diseases, such as thyroid dysfunction and adrenal dysfunction, chronic metabolism disorder, chronic renal failure including nephrosis, uremia, cancer, dyscrasia, adverse effects of antitumor agent, gastrointestinal(including hepatic and pancreatic) impairment and so on.

Background of the Invention

More than 60% of the cause of Japanese death is adult disease, such as hypertension, arteriosclerosis, diabetes and so on. Generally, adult disease is caused mainly by deterioration of endogenous factor, such as endocrinological function and/or lipid metabolism function, whose controlling mechanism is not yet known in many aspects. Category of therapeutic agent for treatment of

hyperlipidemia in terms of action mechanism is classified as absorption inhibition of cholesterol or choleic acid, inhibition of cholesterol synthesis and action on very low density protein metabolism (VLDL). Hyperlipidemia accompanied with decrease of high density protein(HDL)-cholesterol is a significant risk factor of arteriosclerotic diseases comprising ischemic heart disease, hypertension, ischemic encephalopathy and so on. It has been found that pharmacological effect of a therapeutic agent for treatment of hyperlipidemia is not only decreasing serum lipid, but also improving lowered HDL-cholesterol which is a progressive factor of hyperlipidemia and/or lowered lecithin-cholesterol acyltransferase (LCAT) so as to prevent from progression of hyperlipidemia to arteriosclerotic disease by removing stagnation of metabolism.

However, a pharmaceutical agent for improving lipid metabolism which improves lipid metabolism pertinently is not yet developed. As a therapeutic agent for treatment of abnormal lipid metabolism which is used clinically at present, Pravastatin sodium salt which inhibits 3-hydroxy-3-methyl-glutaryl-CoA reducing enzyme or clofibrate having clarification action on serum lipid is exemplified for hyperlipidemia and alimentation including high calory infusion is exemplified for hypolipidemia.

Various therapeutic agents are used depending on symptoms regardless of primary disease. Thus, there has not been any pharmaceutical agent for improving and/or treating abnormal lipid metabolism caused by imbalance of endocrinological dysfunction

associated with lipid and carbohydrate metabolism. As the results of various causes, diabetes shows pathological features which are induced by imbalance of insulin and glucagon and endocrinological dysfunction-including pancreatic dysfunction, leading to metabolic disorder of liver, lipid, muscle, skin, kidney and so on. A role of adipose tissue is to stock as lipid in the case of excess of energy and to decompose lipid and supply to living body as an energy source if required. In the case of diabetes, synthesis is inhibited in adipose tissue and decomposition is stimulated because of insulin deficiency and excessive glucagon. That is, by decrease of insulin action, sucrose intake decrease in adipose tissue, resulting in decrease of production of α -glycerophosphoric acid, NADH and NADPH and decrease of synthesis of fatty acid and reesterification. On the other hand, insulin deficiency and excessive glucagon activates hormone-sensitive lipase to decompose lipid. Mechanism of development of fatty liver is thought that impairment occurs in either process of triglyceride synthesis or synthesis and secretion of VLDL in liver.

Stimulation of fatty acid synthesis in liver, increase of fatty acid mobilization from peripheral adipose tissue to liver, decrease of ability to oxidize fatty acid and decrease of synthesis and secretion of lipoprotein in liver are thought to be examples of the above mechanism. For example, in fatty liver associated with obesity, mobilization of fatty acid from peripheral adipose tissue increase because of increase of the content of peripheral adipose tissue. Recently, attention is focused on the relation between

development of fatty liver and increase of fatty acid in portal vein due to accumulation visceral lipid. In addition, acidity of acetyl CoA carboxylase which is a rate determining enzyme in de novo synthesis of fatty acid in liver is enhanced in obesity, suggesting enhancement of synthesis of fatty acid in liver. On the other hand, in obesity, hyperinsulinemia is induced to compensate increase of insulin resistance in peripheral adipose tissue.

It is known that fatty liver is accompanied with obesity with hyperinsulinemia at a high rate, comparing to obesity with normal insulin secretion. Thus, with respect to abnormal linked metabolism of carbohydrate and lipid, there has not been any pharmaceutical agent for improving lipid metabolism tactically by stimulating synthesis of serum lipoprotein carrying lipid in blood, improving mobilization thereof to lipid composition and metabolism thereof, enhancing LCAT activity associated with esterification, decreasing free cholesterol level, increasing HDL-cholesterol level, stimulating choleic excretion, decreasing serum lipid level, effluxing triglyceride accumulated in hepatic cell and improving fatty liver.

Considering these situations, the present inventors have eagerly investigated on a therapeutic agent for treatment of abnormal linked metabolism of carbohydrate and lipid and found that TCF-II specifically relates to mobilization of serum lipoprotein carrying lipid in blood to lipid composition and metabolism thereof. Inhibition of synthesis thereof, decrease of LCAT activity

associated with esterification and decrease of HDL accompanied therewith are regarded as serious risk factors of adult diseases accompanied with disorder of lipid metabolism such as hyperlipidemia, arteriosclerosis and so on. The present inventors found that TCF-II significantly enhanced LCAT activity in vivo, improved hypo-high-density-lipoproteinemia, further, stimulated the mobilization by increase serum lipoprotein and normalized serum lipid level. An object of the present invention is to provide a therapeutic agent having efficacy to normalize abnormal lipid metabolism comprising TCF-II as an effective ingredient.

Summary of the Invention

An object of the present invention is to provide a therapeutic agent for treatment of disorder of lipid metabolism comprising TCF-II as an effective ingredient.

Another object is to provide a therapeutic agent for treatment having an action of decreasing high level of serum lipid and normalizing hypolipidemia and having efficacy to improve fatty liver.

Brief Description of Drawings

Figure 1 shows the method of fractionation of serum VLDL fractions in example 4.

Figure 2 shows the method of fractionation of serum HDL fraction and HDL₂ fraction in example 4.

Figure 3 shows concentration of serum triacylglycerol in example 4.

Figure 4 shows concentration of serum ester-type cholesterol and serum free cholesterol and LCAT activity in blood in example 4.

Figure 5 shows administration protocol in example 5.

Figure 6 shows content of triacylglycerol in tissue and concentration of serum phospholipid and serum total cholesterol in example 5.

Figure 7 shows concentration of serum albumin, serum HDL-cholesterol, serum free cholesterol and serum total cholesterol in example 6.

Figure 8 shows administration protocol in example 7.

Figure 9 shows LCAT activity in example 7.

Figure 10 shows administration protocol in example 8.

Figure 11 shows concentration of serum HDL-cholesterol, serum cholesterol and serum total cholesterol.

Detailed Description of the Invention and Preferred Embodiments

A therapeutic agent for treatment of disorder of lipid metabolism comprising TCF-II as an effective ingredient of the present invention is useful for a therapeutic agent for treatment of disorder of lipid metabolism caused by diabetes, endocrinological diseases, such as thyroid dysfunction and adrenal dysfunction, chronic metabolic disorder, chronic renal failure including nephroses, uremia, cancer, dyscrasia, adverted effect of antitumor agent, gastrointestinal (including hepatic and pancreasic) impairment and so on. As a disorder of lipid metabolism described above, hypercholesterolemia, hypolipidemia or hypo-high-density-lipoproteinemia can be exemplified.

TCF-II as an effective ingredient of the present invention is a known protein derived from human fibroblast and characteristics thereof is shown as follows:

i) Molecular weight (SDS electrophoresis)

under non-reducing conditions: $78,000 \pm 2,000$, or $74,000 \pm 2,000$

under reducing conditions : $52,000 \pm 2,000$ (common band)

$30,000 \pm 2,000$ (band B)

$26,000 \pm 2,000$ (band C)

ii) Isoelectric point : 7.4-8.6

iii) Length of amino acid : 723

TCF-II described above can be obtained by concentrating culture

broth of human fibroblast, adsorbing it on ion exchanger and purifying the eluted substance by affinity chromatography (WO90/10651) or by gene manipulation (WO92/01053).

As TCF-II of an effective ingredient of the present invention, TCF-II derived from human fibroblast or TCF-II produced by gene manipulation based on gene sequence described in patent WO90/10651 using microbial or other cell can be used. In addition, TCF-II obtained by gene manipulation described in patent WO92/01053 can be also used. TCF-II with different carbohydrate chain or without carbohydrate chain due to the difference of host cell or microbial organism can be also used in the above case.

However, TCF-II with carbohydrate chain can be preferably used because carbohydrate chain relates to metabolic rate in a living body. TCF-II obtained by these method can be concentrated and purified by usual methods of isolation and purification. For example, precipitation method using organic solvent, salting-out, gel filtration, affinity chromatography using monoclonal antibody and electrophoresis etc. are exemplified. Purification by affinity chromatography using monoclonal antibody can be carried out by using monoclonal antibody described in Japanese published unexamined patent application 97(1993). Obtained TCF-II can be freeze dried or preserved in a freezer. A therapeutic agent for treatment of disorder of lipid and carbohydrate linked metabolism can be administered intravenously, intramuscularly or

subcutaneously as injections. These pharmaceutical preparations can be produced according to known pharmaceutical preparation method. Controlling agent of pH, buffer, stabilizer and so on can be added thereto, if necessary. Administration dosage of the pharmaceutical preparation of the present invention in a patient can be varied depending on symptom, health condition, age or body weight etc. of a patient. As purified TCF-II, not limiting specifically, 0.6mg-600mg/ adult man/day can be administered. Preferably, a pharmaceutical preparation comprising 6mg-60mg of TCF-II can be administered once or more per day.

The present invention will be described in detail by exemplifying examples but the scope of the present invention is not limited to these examples.

[Example 1]

Purification of TCF-II

According to the method described in patent WO90/10651 and the method by Higashio et.al. (Higashio, K. et. al., B.B.R.C., vol.170, pp 397-404 (1990)), purified TCF-II was obtained by cell culture. Human fibroblast cells IMR-90 (ATCC CCL 186) 3×10^6 were inoculated into a roller bottle including 100ml DMEM medium with 5% calf serum and cultured by rolling it at rotation rate of 0.5-2 times/min. for 7 days. When total cell number became 1×10^7 , cells were collected on the bottom surface by trypsin treatment, 100g of 5-9 mesh

sterilized ceramics were added thereto and stationary culture thereof was kept for 24 hours. Then, 500 ml of the above culture broth was added thereto and culture was continued. The whole culture medium was collected every 7-10 days and fresh medium was supplemented. Thus, production was kept for 2 months and 4 L of culture broth per one bottle was collected. Specific activity of the culture broth obtained as described above was 32 μ /ml.

The culture broth (750 L) was concentrated by ultrafiltration using a membrane filter (MW 6,000 cut; Amicon) and purification by serial column chromatography using CM-Sephadex C-50 (Pharmacia), ConA-Sepharose (Pharmacia), Mono S column (Pharmacia) and Heparin-Sepharose (Pharmacia) was carried out to give purified TCF-II.

[Example 2]

Production of Recombinant TCF-II

According to the method described in patent publication W092/01053, cells transformed with TCF-II gene were cultured and purified TCF-II was obtained. Transformed Namalwa cells were cultured and 20 L of culture broth was obtained. This culture broth was purified by CM-Sephade x C-50 (Pharmacia) chromatography, ConA-Sepharose CL-6B (Pharmacia) chromatography and HPLC packed with Mono S (Pharmacia) and about 11 mg of active TCF-II was obtained.

[Example 3]

Production of Pharmaceutical Preparations

Production examples of injections of recombinant TCF-II obtained in the above example 2 are shown below.

1	TCF-II	20 µg
	human serum albumin	100 mg

The above composition was dissolved in 0.01M phosphate buffer solution with pH of 7.0 and the whole volume thereof was made up to 20 ml. After sterilization, the sample was divided into a vial 2ml each and freeze dried, followed by sealing thereof.

2	TCF-II	40 µg
	Tween 80	1 mg
	human serum albumin	100 mg

The above composition was dissolved in a physiological saline solution for injections and the whole volume thereof was made up to 20 ml. After sterilization, the sample was divided into a vial 2ml each and freeze dried, followed by sealing thereof.

3	TCF-II	20 µg
	Tween 80	2 mg
	sorbitol	4 g

The above composition was dissolved in 0.01M PBS with pH of 7.0 and the whole volume thereof was made up to 20ml. After sterilization, the sample was divided into a vial 2ml each and freeze dried,

followed by sealing thereof.

4	TCF-II	40 µg
	Tween 80	1 mg
	glycine	2 g

The above composition was dissolved in a physiological saline solution for injections and the whole volume thereof was made up to 20 ml. After sterilization, the sample was divided into a vial 2 ml each and freeze dried, followed by sealing thereof.

5	TCF-II	40 µg
	Tween 80	1 mg
	sorbitol	2 g
	glycine	1 g

The above composition was dissolved in a physiological saline solution for injections and the whole volume thereof was made up to 20ml. After sterilization, the sample was divided into a vial 2ml each and freeze dried, followed by sealing thereof.

6	TCF-II	20 µg
	sorbitol	4 g
	human serum albumin	50 mg

The above composition was dissolved in 0.01M PBS with pH of 7.0 and the whole volume thereof was made up to 20 ml. After sterilization, the sample was divided into a vial 2ml each and freeze dried, followed by sealing thereof.

7	TCF-II	40 µg
	glycine	2 g
	human serum albumin	50 mg

The above composition was dissolved in a physiological saline solution for injections and the whole volume thereof was made up to 20ml. After sterilization, the sample was divided into a vial 2ml each and freeze dried, followed by sealing thereof.

8	TCF-II	40 µg
	human serum albumin	50 mg

The above composition was dissolved in 0.01M PBS and the whole volume thereof was made up to 20 ml. After sterilization, the sample was divided into a vial 2 ml each and freeze dried, followed by sealing thereof.

[Example 4]

Stimulation of lipoprotein secretion and plasma LCAT activity in Normal Rats by TCF-II

Male wistar rats (8 weeks old, 300g, 5 rats/group) were intravenously injected twice a day with 500 µg/kg of TCF-II (1000 µg/kg/day) for 4 days. Rats of control group were injected with only vehicle. Twelve hours after the last injection, blood was taken from posterior cava under ether anesthesia. Sera and plasma were collected for following analysis of lipid in lipoprotein and

LCAT activity, respectively. For analysis of lipoprotein, VLDL was separated according to the methods of Wilson et. al. (Clin. Chem., vol.20, 394-395, 1974), and HDL₂ and HDL₃ were separated according to the methods of Gidez et. al. (J. Lipid Res., vol.23, 1206-1223, 1982). Schematic representations of the above procedures are illustrated in figure 1 and 2. The concentrations of triacylglycerol, total cholesterol and free cholesterol were determined enzymatically with the commercially available kits (Wako Pure Chemical Industries, Osaka, Japan). The concentration of esterified cholesterol was calculated according to the following relationship, {[concentration of total cholesterol] - [concentration of free cholesterol]}. Plasma LCAT activity was determined according to the methods of Glomset-Wright (Methods in Enzymology, vol.15, 543, 1969).

The concentration of triacylglycerol in VLDL was significantly higher in TCF-II-injected rats than in control rats (Figure 3). This result suggested that TCF-II stimulates secretion of VLDL and release of triacylglycerol into blood from liver tissue. In addition, plasma LCAT activity was significantly enhanced and the concentration of HDL cholesterol, especially esterified cholesterol, markedly increased in TCF-II-injected rats (Figure 4), which showed that TCF-II stimulated secretion of LCAT from liver so as to produce HDL.

[Example 5]

Reduction of Highly Accumulated Liver Triacylglycerol in

Alcohol-induced Fatty Liver Rats by TCF-II

Liquid diet was prepared according to the method of Liber, et al (Trans. Assoc. Am. Physician. vol.76, 289-301, 1963). Male wistar rats (7 weeks old, 280g, 10-13 rats/group) were fed with the liquid diet containing 5% alcohol for 5 weeks and induced to fatty liver. Control rats were fed with the liquid control diet of equal calories. Fatty liver rats were intravenously injected twice a day with 50 and 500 $\mu\text{g/kg}$ of TCF-II (100 and 1000 $\mu\text{g/kg/day}$, respectively) for the last 7 days of alcohol feedings. Rats of control group were injected with only vehicle.

Twelve hours after the last TCF-II injection, the animals were sacrificed and livers were excised. The liver lipids were extracted by 30 volumes of chloroform and methanol (2:1), and the contents of triacylglycerol, phospholipid and total cholesterol were determined with the commercially available kits (Wako Pure Chemical Industries, Osaka, Japan). The administration protocol is shown in figure 5.

The highly accumulated liver triacylglycerol in fatty liver rats were dose-dependently diminished by TCF-II injection. The liver phospholipid and cholesterol levels, however, remained unchanged (Figure 6). These results also suggested that TCF-II increases triacylglycerol secretion from liver and improves fatty liver.

[Example 6]

Reduction of High Serum Cholesterol Concentration in

α -naphthylisothiocyanate (ANIT)-induced Cholestasis Rats by TCF-II

Male wistar rats (7 weeks old, 280g, 10 rats/group) were orally administered 100 mg/kg of ANIT for 5 weeks and induced to cholestasis. Cholestasis rats were intravenously injected twice a day with 50, 150, 500 and 1500 μ g/kg of TCF-II (100, 300, 1000 and 3000 μ g/kg/day, respectively) for 3 days immediately after administration of ANIT. Twelve hours after the last injection, blood was taken from posterior cava under ether anesthesia. Sera were collected for following measurement of serum lipid and albumin concentration. The serum concentrations of triacylglycerol, free cholesterol, total cholesterol and albumin were determined by Hitachi 7150 Autoanalyzer.

The serum concentrations of triacylglycerol, free cholesterol, total cholesterol of the cholestasis rats was significantly higher than those of normal rats. TCF-II injection reduced the serum lipid concentrations and increased the serum albumin concentration in a dose-dependent manner (Figure 7). From these results, TCF-II improves hyperlipidemia accompanied by cholestasis.

[Example 7]

Stimulation of plasma LCAT activity in Dogs by TCF-II

Female and male Beagles (1-5 years old, about 10 kg, 5 dogs/group) were intravenously injected twice a day with 50, 150 and 500 μ g/kg of TCF-II (100, 300 and 1000 μ g/kg/day, respectively) for 7 days.

Dogs of control group were injected with only vehicle. Before and during TCF-II injection, blood was taken from forelimb vein and plasma were collected for following analysis of LCAT activity. The administration protocol is shown in figure 8.

Plasma LCAT activity was determined according to the methods of Glomset-Wright (Methods in Enzymology, vol.15, 543, 1969). Plasma LCAT activities of the TCF-II injected animals markedly increased (Figure 9), which showed that TCF-II stimulates secretion of LCAT from liver and lipid metabolism in normal dogs.

[Example 8]

Elevation of Low Serum Cholesterol, Glucose Concentration and Low Liver Glycogen contents in Dimethylnitrosamine (DEM)-induced Cirrhosis Rats by TCF-II

Male wistar rats (8 weeks old, 300g, 8-10 rats/group) were intraperitoneally administered 3 times a week 10 µg/kg of DEM for 4 weeks and induced to cirrhosis.

Cirrhosis rats were intravenously injected twice a day with 5, 50 and 100 µg/kg of TCF-II (10, 100 and 1000 µg/kg/day, respectively) for 4 weeks during administration of DEM. The administration protocol is shown in figure 10. Twelve hours after the last injection, blood was taken from posterior cava under ether anesthesia. Sera were collected for following measurement of serum cholesterol concentration. The serum concentrations of HDL-cholesterol, free cholesterol and total cholesterol were

determined by Hitachi 7150 Autoanalyzer. The serum concentrations of esterified cholesterol were calculated according to the relationship described in Example 4.

The serum concentrations of HDL-cholesterol and esterified cholesterol of the cirrhosis rats were significantly lower than those of normal rats, but the concentration of free cholesterol higher. TCF-II injection elevated the serum HDL-cholesterol and esterified cholesterol concentrations and reduced the serum free cholesterol concentration significantly (Figure 11). From these results, TCF-II improves hypo-high-density-lipoproteinemia accompanied by cirrhosis.

Industrial Applicability


The present invention is to provide a therapeutic agent for treatment of disorder of lipid metabolism comprising TCF-II as an effective ingredient. The therapeutic agent of the present invention normalizes hyperlipidemia and hypolipidemia. It also improves fatty liver.

Claims

1. A therapeutic agent for treatment of disorder of lipid metabolism comprising TCF-II as an effective ingredient.
2. The therapeutic agent for treatment of disorder of lipid metabolism according to claim 1, wherein disorder of lipid metabolism is hypercholesterolemia.
3. The therapeutic agent for treatment of disorder of lipid metabolism according to claim 1, wherein disorder of lipid metabolism is hypolipidemia.
4. The therapeutic agent for treatment of disorder of lipid metabolism according to claim 1, wherein disorder of lipid metabolism is hypo-high-density-lipoproteinemia.
5. The therapeutic agent for treatment of disorder of lipid metabolism according to claim 1, wherein disorder of lipid metabolism is fatty liver.

Fig. 1


1/11



Formula for calculating lipid concentration of VLDL fraction

$$[\text{VLDL}] = [\text{Serum}] - [\text{A}]$$

Fig. 2



Formula for calculating lipid concentration of each lipid fraction

$$[LDL] = [A] - [VLDL]$$

$$[HDL_2] = [B] - [C]$$

$$[HDL_3] = [C]$$

Fig. 3

3/11





Fig. 4


4/11



5/11

Fig. 5

Administration protocol





Duration of feeding alcoholic diet and control diet

Duration of TCF-II administration

6/11


Fig. 6



means \pm standard error
(n=5) **: p < 0.01


7/11

Fig. 7




8/11

Fig. 8

Administration protocol


9/11

Fig. 9




10/11

Fig.10



11/11

Fig. 11



INTERNATIONAL SEARCH REPORT

International Application No.
PCT/JP 95/02707A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K38/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 588 477 (SNOW BRAND MILK PRODUCTS CO., LTD.) 23 March 1994 see the whole document ---	1-5
X	EP,A,0 462 277 (SNOW BRAND MILK PRODUCTS CO., LTD.) 27 December 1991 cited in the application see the whole document -----	1-5

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- * "A" document defining the general state of the art which is not considered to be of particular relevance
- * "E" earlier document but published on or after the international filing date
- * "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- * "O" document referring to an oral disclosure, use, exhibition or other means
- * "P" document published prior to the international filing date but later than the priority date claimed

* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* "&" document member of the same patent family

Date of the actual completion of the international search

15 April 1996

Date of mailing of the international search report

23.04.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patendaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Moreau, J

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/JP 95/02707

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-588477	23-03-94	JP-A- 6040934	15-02-94
		JP-A- 6040935	15-02-94
		JP-A- 6056692	01-03-94
		AU-B- 4194593	20-01-94
		CA-A- 2100720	17-01-94
		ZA-A- 9305119	24-02-94

EP-A-462277	27-12-91	DE-D- 69019962	13-07-95
		DE-T- 69019962	23-11-95
		JP-B- 7068272	26-07-95
		AT-T- 123498	15-06-95
		AU-B- 629314	01-10-92
		AU-B- 5177490	09-10-90
		CA-A- 2049017	11-09-90
		WO-A- 9010651	20-09-90
